

Surface abundances of OC supergiants[★] (Research Note)

F. Martins¹, S. Foschino¹, J.-C. Bouret², R. Barbá³, and I. Howarth⁴

¹ LUPM, Université de Montpellier, CNRS, Place Eugène Bataillon, 34095 Montpellier, France
e-mail: fabrice.martins@umontpellier.fr

² Aix Marseille Université, CNRS, LAM (Laboratoire d’Astrophysique de Marseille) UMR 7326, 13388 Marseille, France

³ Departamento de Física y Astronomía, Universidad de La Serena, Cisternas 1200 N, La Serena, Chile

⁴ Department of Physics & Astronomy, University College London, Gower St, London WC1E 6BT, UK

Received 10 February 2016 / Accepted 26 February 2016

ABSTRACT

Context. Some O and B stars show unusually strong or weak lines of carbon and/or nitrogen. These objects are classified as OBN or OBC stars. It has recently been shown that nitrogen enrichment and carbon depletion are the most likely explanations for the existence of the ON class.

Aims. We investigate OC stars (all being supergiants) to check that surface abundances are responsible for the observed anomalous line strengths.

Methods. We perform a spectroscopic analysis of three OC supergiants using atmosphere models. A fourth star was previously studied by us. Our sample thus comprises all OC stars known to date in the Galaxy. We determine the stellar parameters and He, C, N, and O surface abundances.

Results. We show that all stars have effective temperatures and surface gravities fully consistent with morphologically normal O supergiants. However, OC stars show little, if any, nitrogen enrichment and carbon surface abundances consistent with the initial composition. OC supergiants are thus barely chemically evolved, unlike morphologically normal O supergiants.

Key words. stars: massive – stars: abundances – stars: fundamental parameters – stars: early-type

1. Introduction

Some OB stars show lines of carbon and nitrogen with unusual morphologies. Jaschek & Jaschek (1967) were the first to report very weak nitrogen optical lines in two supergiants. Subsequent observations of OB stars revealed a number of objects that had similar spectral properties and also showed strong carbon lines (Walborn 1970; Jaschek & Jaschek 1974). Some OB stars displayed strong nitrogen lines and weak carbon lines, too. Walborn (1971) introduced the spectral types OBN and OBC, by analogy with Wolf-Rayet stars where similar line characteristics are seen (see also Walborn 1976). The number of OBN/OBC stars is small: according to the summary of Walborn et al. (2011), only thirteen (four) Galactic O stars have an ON (OC) spectral type.

In Wolf-Rayet stars, the strength of carbon and nitrogen lines is related to their evolutionary status. Winds have removed the external layers and exposed inner regions where H and He burning products are present. Walborn (1976) lists the possible mechanisms explaining the appearance of OBN/OBC stars. Chemical enrichment/depletion is a plausible explanation. Recently, Martins et al. (2015b) have quantitatively studied the surface abundances of ON stars¹. They concluded that strong nitrogen enrichment and carbon depletion was the reason for their spectral morphology. The physical origin of this chemical pattern remains unclear though.

In this follow-up study, we focus on OC stars. Martins et al. (2015a) analyzed a large sample of O stars including one OC object (HD 152249). The latter turned out to show little nitrogen enrichment, unlike similar objects without the “C” qualifier. Here, we study the three additional OC stars of the sample of Walborn et al. (2011).

2. Observations and sample

The spectroscopic data for HD 104565, HD 152424, and HD 154811 were obtained in the context of the OWN project (Barbá et al. 2010, 2014) conducted at ESO/La Silla with the FEROS instrument mounted on the 2.2 m Max-Planck telescope (Table 1). They have a resolving power of 48 000. They were reduced by the fully automated pipeline distributed by ESO. The spectrum of HD 152249 was acquired with the ESPaDOnS spectropolarimeter mounted on the Canada–France–Hawaii Telescope. Its analysis was presented in Martins et al. (2015a), and we adopt the parameters obtained in this study.

3. Modeling and spectroscopic analysis

We used the atmosphere code CMFGEN Hillier & Miller (1998) to analyze the surface properties of the OC stars. The method we used is the same as that presented in Martins et al. (2015b), and we refer the reader to this paper for a full description. CMGFEN computed non-LTE, spherically extended atmosphere models that include line-blanketing. Synthetic spectra calculated from

[★] Based on observations obtained at the ESO/La Silla Observatory under programs 081.D-2008, 083.D-0589, 089.D-0975.

¹ A summary of previous quantitative studies of ON stars can be found in Martins et al. (2015b).

Table 1. Observational information.

Star	Sp.T. ¹	Instrument	Date of observation
HD 104565	OC9.7 Iab	FEROS	12 May 2008
HD 152249 ²	OC9 Iab	ESPaDOnS	14 Jul. 2011
HD 152424	OC9.2 Ia	FEROS	05 Apr. 2015
HD 154811	OC9.7 Ib	FEROS	average

Notes. 1– Spectral types are from Sota et al. (2011, 2014). 2– Star analyzed by Martins et al. (2015a).

1 the atmosphere models are compared to observations to determine the stellar parameters. The main parameters were obtained as follows:

- 4 – *Rotation and macroturbulence:* we used the Fourier-transform of O III $\lambda 592$ to determine the projected equatorial rotational velocity $V \sin i$ (Gray 1976). We then convolved a synthetic spectrum (with T_{eff} and $\log g$ estimated from spectral type) with a radial–tangential profile. The comparison of this synthetic spectrum with the observed profile yielded the macroturbulent velocity v_{mac} .
- 11 – *Effective temperature:* the ionization-balance method was used to constrain T_{eff} , taking advantage of the numerous He I and He II lines present in the optical spectra. The ionization balance of carbon (based on C III lines and C II 4267) is consistent – within the error bars – with the effective temperature determined from helium lines.
- 17 – *Surface gravity:* $\log g$ was determined from the width of the Balmer lines wings. Broader wings are observed for higher surface gravities.

20 Once these parameters were constrained, we ran models with different surface abundances (for He, C, N, and O). For each star and each element, a set of clean lines was selected from the observed spectrum. These lines were quantitatively compared to the synthetic spectra by means of a χ^2 analysis from which we derived the surface abundance and associated uncertainty (see Martins et al. 2015a,b). Abundance determinations were performed assuming a microturbulent velocity of 10 km s^{-1} .

28 4. Results

29 The best fits to the optical spectra of the target stars are presented in Appendix A. The stellar parameters and surface abundances are summarized in Table 2. For all stars, we found that He/H = 0.1 led to good fits (Figs. A.1 to A.3). Figure 1 shows the $\log g - T_{\text{eff}}$ diagram with evolutionary tracks from Ekström et al. (2012). We added the morphologically normal late-O supergiants of Martins et al. (2015a) for comparison. All stars are located at, or just beyond, the terminal age main sequence. They are relatively clustered with initial masses between ~ 20 and $\sim 40 M_{\odot}$. OC and normal O supergiants are located at the same position of the diagram. As a result, OC stars do not have peculiar effective temperatures or surface gravities.

41 Figure 2 shows the ratio of nitrogen-to-carbon abundance as a function of surface gravity. The evolutionary tracks of Ekström et al. (2012) for the range of masses deduced from Fig. 1 are overplotted. The difference between normal and OC stars is striking. While normal O supergiants are accounted for relatively well by the rotating-star evolutionary tracks, the OC stars have N/C ratios lower by 0.6–0.8 dex (factor 4 to 6). In fact, OC stars are almost consistent with no chemical processing (N/C 0.0 to 0.3 dex higher than the initial value). Figure 3 shows the position




Fig. 1. $\log g - \log(T_{\text{eff}})$ diagram for the sample stars (filled circles) and comparison late-type supergiants (O9–9.7I – open circles). Typical uncertainties are shown in the upper-left corner. Evolutionary tracks from Ekström et al. (2012) including rotation are overplotted, labeled by initial masses.




Fig. 2. $\log(N/C) - \log g$ diagram for the sample stars (filled circles) and comparison late-type supergiants (open circles). Evolutionary tracks including rotation from Ekström et al. (2012) are overplotted, labeled by initial masses. The dotted line shows the initial chemical composition. For non-rotating models, N/C remains at the initial value (dotted line) throughout the main sequence and early post-main sequence phases.

of the OC stars (and comparison supergiants) in the $\log(N/C) - \log(N/O)$ diagram. OC stars are fully consistent with stars having experienced very little processing through the CN or CNO cycles. Morphologically normal supergiants are more chemically evolved.

Table 2. Parameters of the sample stars.

Star	Spectral type	T_{eff} [kK]	$\log g$	$\log g_c$	$V \sin i$ [km s $^{-1}$]	v_{mac} [km s $^{-1}$]	C/H [10^{-4}]	N/H [10^{-4}]	O/H [10^{-4}]
HD 104565	OC9.7 Iab	28	3.00	3.01	60	90	>2.0	$2.0^{+0.7}_{-0.7}$	>1.7
HD 152249	OC9 Iab	31	3.25	3.25	43	48	$2.8^{+0.4}_{-0.4}$	$1.3^{+0.7}_{-0.7}$	>7.0
HD 152424	OC9.2 Ia	29	3.10	3.11	60	90	$2.0^{+1.4}_{-0.8}$	$0.9^{+0.2}_{-0.4}$	>2.0
HD 154811	OC9.7 Ib	28	3.10	3.12	110	65	$2.0^{+0.8}_{-0.5}$	$1.2^{+1.0}_{-0.6}$	$4.6^{+1.4}_{-1.1}$

Notes. Uncertainties on T_{eff} , $\log g$, $V \sin i$, and v_{mac} are ~ 1.5 kK, 0.15 dex, 10, and 20 km s $^{-1}$ respectively. $\log g_c$ is the surface gravity corrected for centrifugal acceleration. Abundances are number ratios. Values for HD 152249 are from Martins et al. (2015a).




Fig. 3. $\log(N/C) - \log(N/O)$ diagram for the sample stars (filled circles) and comparison late-type supergiants (open circles) from Martins et al. (2015a). Solid lines indicate the prediction of nucleosynthesis through the partial CN or complete CNO cycle.

5. Discussion

Figure 2 reveals that OC supergiants are much less chemically evolved than morphologically normal O supergiants. They are also less evolved than expected for stars of initial mass 20–40 M_{\odot} , assuming a standard initial rotational velocity of ~ 300 km s $^{-1}$. Indeed, the evolutionary tracks of Ekström et al. (2012) with this initial rotation predict values of $\log(N/C)$ between 0.6 and 1.0, while OC supergiants have $\log(N/C)$ around -0.3. It is conceivable that OC supergiants are the descendants of stars rotating very slowly on the main sequence. In that case, single-star evolution predicts that N/C should remain constant down to $\log g \sim 0.0$, corresponding to the red supergiant phase. In Fig. 2, such an evolution would be represented by the dotted line. If this explanation holds, OC supergiants would represent the descendants of main-sequence stars at the low-velocity tail of the initial rotational velocity distribution. Since the latter is continuous (Ramírez-Agudelo et al. 2013), one would expect to find O supergiants with $\log(N/C)$ over the full range -0.3–1.0 in the $\log(N/C) - \log g$ diagram (Fig. 9 of Brott et al. 2011). In Fig. 2, there seems to be a deficit of stars for $\log(N/C)$ between -0.2 and 0.2 dex. However, the number of stars in this diagram remains small. Interestingly, Hillier et al. (2003) studied a O7

supergiant and a OC7.5 giant in the SMC. From the stellar properties and surface abundances, they concluded that the latter was a slow rotator that did not experience strong mixing.

If slow rotation is the explanation for the chemical appearance of OC stars, their initial masses should be slightly higher than those of morphologically normal O supergiants. Indeed, evolutionary tracks without rotation are shifted to the right in Fig. 1 compared to tracks including rotation. Consequently, higher masses are needed to reproduce the position of OC stars (40–60 M_{\odot} instead of 20–40 M_{\odot}). OC stars may thus differ from morphologically normal stars in that respect.

Slow rotation does not explain all the properties of OC stars. In particular, HD 154811 has a present-day rotational velocity of 110 km s $^{-1}$. This value is relatively large for O supergiant: Simón-Díaz & Herrero (2014) report $V \sin i$ for O9-9.7 supergiants in the range 48–132 km s $^{-1}$, with an average value of 88 ± 30 km s $^{-1}$. Given that the projected rotational velocity drops significantly when a star leaves the main sequence, one would expect HD 154811 to have an initial $V \sin i$ larger than 110 km s $^{-1}$. For instance under the assumption of angular momentum conservation and solid-body rotation, the ratio of initial to present $V \sin i$ is equal to the ratio of present to initial stellar radius. Assuming a factor of three for the latter ratio (according to the tracks of Ekström et al. 2012), the initial rotational velocity of HD 154811 would have been 330 km s $^{-1}$, a fairly high value. For the other targets, the initial projected rotational velocity would have been 120–160 km s $^{-1}$.

The OBN/OBC classification was introduced by analogy to Wolf-Rayet stars (Walborn 1971). In WC stars, the appearance of strong lines of carbon are caused by the large surface content of carbon produced by helium burning. One could speculate that the same is true for OC stars. However, in that case there should be no hydrogen at the surface of the stars (Crowther 2007). In addition, helium should be the dominant element, which is clearly not the case. As a result, the abundance patterns of OC stars are not due to helium-burning products.

Surface abundances can be altered by mass transfer in binary systems. Bolton & Rogers (1978) investigated binarity among OC stars. Their results indicated a low binary fraction, although results were not conclusive for some stars. Levato et al. (1988) revisited this question and reached more robust conclusions: OC stars usually do not show the radial velocity variations typical of spectroscopic binaries. Sana et al. (2008) obtained time series spectroscopy of HD 152249 and did not detect any sign of a companion. The interferometric search for companions around O stars by Sana et al. (2014) confirms that HD 152249, HD 152424, and HD 154811 do not have bright companions. The ongoing OWN spectroscopic survey of O stars confirm these results: only HD 152424 shows radial velocity

1 variations greater than 5 km s^{-1} . HD 152424 may be a SB1 with
 2 a low mass component (Barbá et al., in prep). All other OC stars
 3 are probably single objects. Consequently, the chemical abundances
 4 of OC stars are unlikely to be due to binary interaction.
 5 Besides, if mass transfer can explain nitrogen enrichment and
 6 carbon depletion, it cannot explain that surface abundances are
 7 almost unaltered. This would have to be due to tidal interaction
 8 that would suppress internal mixing processes.

9 Interestingly, Bouret et al. (2008) report very little nitrogen
 10 enrichment and almost solar carbon abundance in the star ζ Ori.
 11 Sota et al. (2014) classify it as “Nwk”, meaning that the nitrogen
 12 lines are weak, as in OC stars. And ζ Ori is known to host a weak
 13 magnetic field (Bouret et al. 2008; Blazère et al. 2015), which is
 14 quite different from the strong dipoles usually observed in mag-
 15 netic O stars (e.g., Wade & MiMeS Collaboration 2015). Martins
 16 et al. (2015a) showed that the chemical properties of O stars with
 17 dipole fields did not significantly differ from stars without mag-
 18 netic fields. One may wonder whether for the case of a weak,
 19 small scale magnetic field, the situation is different. Meynet et al.
 20 (2011) show that massive stars with magnetic field of only a few
 21 Gauss may slow down very fast provided they are in solid-body
 22 rotation. In that case, there is no time for significant mixing, and
 23 stars appear chemically unevolved (see Fig. 1 of Meynet et al.
 24 2011). A deep spectrolarimetric study of OC stars is needed to
 25 investigate the role of magnetism.

26 All OC stars in the Galaxy are late-type supergiants. Since
 27 the OC phenomenon is due to surface abundances, one can under-
 28 stand that it is best seen in supergiants for which chemical
 29 processing is already strong in morphologically normal stars.
 30 The surface abundances of OC supergiants are similar to those
 31 of O-normal dwarfs, probably because they are relatively chemi-
 32 cally unevolved. Thus, the differential effect leading to the clas-
 33 sification as OC (weak nitrogen lines/strong carbon lines) does
 34 not exist among dwarfs that are only slightly chemically evolved
 35 (Martins et al. (2015a). OC stars may be seen among giants
 36 (Hillier et al. 2003). A recent study by Evans et al. (2015) as-
 37 signs a spectral type OC to two early type O giants in the LMC,
 38 based on the weakness of nitrogen lines.

39 Regarding spectral type, there is a priori no reason for the
 40 OC phenomenon to be restricted to late-type O stars. Walborn
 41 et al. (2010) introduced the “Ofc” classification to characterize
 42 O stars with C III 4650 of the same strength as N III 4640. These
 43 features are seen in emission in Of stars, but the latter is usually
 44 stronger than the former. The distribution of Ofc stars is peaked
 45 at spectral type O5. The strong C III 4650 emission may be due
 46 to a larger carbon content, as in OC stars. However, Martins
 47 & Hillier (2012) pointed out that the formation of C III 4650
 48 in O-type stars depends on several physical parameters (winds,
 49 metallicity, microturbulence, atomic data). A detailed analysis of
 50 Ofc stars using other carbon lines is therefore necessary to check
 51 whether OC and Ofc stars belong to the same class of chemically
 52 unevolved objects.

6. Conclusion

We performed a spectroscopic analysis of three OC supergiants
 using atmosphere models (computed with the code CMFGEN)
 and high-resolution optical spectroscopy. A fourth star was pre-
 viously studied in Martins et al. (2015a). We determined the
 stellar parameters and surface abundances. We compared their
 properties to that of morphologically normal O supergiants. OC
 supergiants have stellar parameters that are fully consistent with
 normal O supergiants. OC stars show little or no nitrogen enrich-
 ment, as well as close-to-initial carbon and oxygen abundances.
 OC supergiants are barely chemically evolved, unlike morpho-
 logically normal O supergiants. Slow rotation may be an expla-
 nation for these surface chemical patterns, although problems
 remain.

Acknowledgements. We thank an anonymous referee for a prompt and construc-
 tive report. We thank John Hillier for making CMFGEN available to the com-
 munity. We acknowledge comments from N.R. Walborn on an early version of
 the paper. R.B. acknowledges support from FONDECYT Project No 1140076.

References

- Barbá, R. H., Gamen, R., Arias, J. I., et al. 2010, in *Rev. Mex. Astron. Astrofis.*
 Conf. Ser., 38, 30
- Barbá, R., Gamen, R., Arias, J. I., et al. 2014, in *Rev. Mex. Astron. Astrofis.*
 Conf. Ser., 44, 148
- Blazère, A., Neiner, C., Tkachenko, A., Bouret, J.-C., & Rivinius, T. 2015, *A&A*,
 582, A110
- Bolton, C. T., & Rogers, G. L. 1978, *ApJ*, 222, 234
- Bouret, J.-C., Donati, J.-F., Martins, F., et al. 2008, *MNRAS*, 389, 75
- Brott, I., de Mink, S. E., Cantiello, M., et al. 2011, *A&A*, 530, A115
- Crowther, P. A. 2007, *ARA&A*, 45, 177
- Ekström, S., Georgy, C., Eggenberger, P., et al. 2012, *A&A*, 537, A146
- Evans, C. J., van Loon, J. T., Hainich, R., & Bailey, M. 2015, *A&A*, 584, A5
- Gray, D. F. 1976, *The observation and analysis of stellar photospheres* (New
 York: Wiley-Interscience), 484
- Hillier, D. J., & Miller, D. L. 1998, *ApJ*, 496, 407
- Hillier, D. J., Lanz, T., Heap, S. R., et al. 2003, *ApJ*, 588, 1039
- Jaschek, M., & Jaschek, C. 1967, *ApJ*, 150, 355
- Jaschek, M., & Jaschek, C. 1974, *A&A*, 36, 401
- Levato, H., Morrell, N., Garcia, B., & Malaroda, S. 1988, *ApJS*, 68, 319
- Martins, F., & Hillier, D. J. 2012, *A&A*, 545, A95
- Martins, F., Hervé, A., Bouret, J.-C., et al. 2015a, *A&A*, 575, A34
- Martins, F., Simón-Díaz, S., Palacios, A., et al. 2015b, *A&A*, 578, A109
- Meynet, G., Eggenberger, P., & Maeder, A. 2011, *A&A*, 525, L11
- Ramírez-Agudelo, O. H., Simón-Díaz, S., Sana, H., et al. 2013, *A&A*, 560, A29
- Sana, H., Gosset, E., Nazé, Y., Rauw, G., & Linder, N. 2008, *MNRAS*, 386, 447
- Sana, H., Le Bouquin, J.-B., Lacour, S., et al. 2014, *ApJS*, 215, 15
- Simón-Díaz, S., & Herrero, A. 2014, *A&A*, 562, A135
- Sota, A., Maíz Apellániz, J., Walborn, N. R., et al. 2011, *ApJS*, 193, 24
- Sota, A., Maíz Apellániz, J., Morrell, N. I., et al. 2014, *ApJS*, 211, 10
- Wade, G. A., & MiMeS Collaboration. 2015, in *Physics and Evolution of*
 Magnetic and Related Stars, eds. Y. Y. Balega, I. I. Romanyuk, & D. O.
 Kudryavtsev, ASP Conf. Ser., 494, 30
- Walborn, N. R. 1970, *ApJ*, 161, L149
- Walborn, N. R. 1971, *ApJ*, 164, L67
- Walborn, N. R. 1976, *ApJ*, 205, 419
- Walborn, N. R., Sota, A., Maíz Apellániz, J., et al. 2010, *ApJ*, 711, L143
- Walborn, N. R., Maíz Apellániz, J., Sota, A., et al. 2011, *AJ*, 142, 150

1 **Appendix A: Best fits to the observed spectra**

2 The fits to the observed spectra are usually of very good qual-
 3 ity. The C III line complex around 4650 Å is not reproduced
 4 in HD 104565 and HD 152424. As stressed by Martins &
 5 Hillier (2012), this line complex depends on details of the mod-
 6 eling (metallicity, blanketing, microturbuence, winds, etc.) and
 7 should not be taken into account for abundance determinations.
 8 Si IV 4089–4116 is also not correctly reproduced by our models
 9 in the same two objects. We stress that we did not try to achieve
 10 a good fit for these lines, which are often a problem for O stars.
 11 A detailed study of their formation processes is needed to under-
 12 stand the present discrepancies.




Fig. A.1. Best fit (red) of the observed spectrum (black) of HD 104565.




Fig. A.2. Best fit (red) of the observed spectrum (black) of HD 152424.




Fig. A.3. Best fit (red) of the observed spectrum (black) of HD 154811.